arXiv:0802.1085v2 [math.RA] 11 Mar 2008

A Note on Auslander Bounds

Jiaqun Wei*
Department of Mathematics, Nanjing Normal University
Nanjing 210097, P.R.China
Email: weijiaqun@njnu.edu.cn

Abstract

Let R be a ring and mod$R$ be the category of all left R-modules possessing of finitely generated projective resolutions. Let $\mathrm{lAb}|_{\mathrm{mod}R}M$ denote the minimal nonnegative integer m (or ∞ if no such integer exists) such that $\mathrm{Ext}_R^i(M,N)=0$ for all $i>m$ and all $N\in \mathrm{mod}R$ satisfying $\mathrm{Ext}_R^i(M,N)=0$ for all sufficiently large i. Auslander once conjectured for any artin algebra R that $\mathrm{lAb}|_{\mathrm{mod}R}M<\infty$ for every $M\in \mathrm{mod}R$, but it is now known that the Auslander Conjecture fails by a counterexample due to Jorgensen and Sega. We show that if $M\in \mathrm{mod}R$ and $\mathrm{Ext}_R^i(M,M\oplus R)=0$ for all sufficiently large i, then the projective dimension of M coincides with $\mathrm{lAb}|_{\mathrm{mod}R}M$. The result is clearly relational to the Auslander-Reitein Conjecture which states that that if $M\in \mathrm{mod}R$ and $\mathrm{Ext}_R^i(M,M\oplus R)=0$ for all $i>0$ then M is projective. As corollaries, we obtain that if R satisfies the condition that $\mathrm{lAb}|_{\mathrm{mod}R}M<\infty$ for every $M\in \mathrm{mod}R$, then an equivalent version of the Wakamatsu-tilting Conjecture, particularly the Auslander-Reitein Conjecture, holds for R, i.e., an R-module $T\in \mathrm{mod}R$ is tilting if and only if (1) $\mathrm{Ext}_R^i(T,T)=0$ for all $i>0$ and (2) there is an exact sequence $0\to R\to T_0\cdots\to T_n\to 0$ for some n, with each $T_i\in \mathrm{add}T$. We prove that all projective ultimately closed artin algebras have the property. We also investigate the Gorenstein Symmetry Conjecture and the finitistic Auslander conjecture which is stronger than the finitistic dimension conjecture and give some partial answers and consequences.

Keywords: Auslander bound Homological conjectures
MSC 2000: 16E10, 16E30, 16E65.

1 Introduction

Throughout this paper, rings are associative with nonzero identities and modules are left unless pointed explicitly. Let R be a ring, we denote by Mod$R$ the category of all left R-modules. The notion mod$R$ denotes the full subcategory of all R-modules possessing of

---

*Supported by the National Natural Science Foundation of China (Grant No. 10601024)

finitely generated projective resolutions. R^o denotes the opposite ring and an R^o-module M means the right R-module M_R.

Auslander once posed the following conjecture (cf. [8]).

(AC) *Let* R *be an artin algebra. Then for every* $M \in \mathrm{mod}R$*, there exists an integer* b_M *such that* $\mathrm{Ext}^i_R(M, N) = 0$ *for all* $i > b_M$ *and for every* R*-module* $N \in \mathrm{mod}R$ *satisfying* $\mathrm{Ext}^i_R(M, N) = 0$ *for all sufficiently large* i*.*

It was known that the Auslander Conjecture holds for all finite dimensional algebras implies that the little finitistic projective dimension conjecture is true for all finite dimensional algebras [8]. However, the Auslander Conjecture fails in general by counterexamples in [11, 18]. Rings satisfying the Auslander conjecture are studied in [4, 9, 16, etc.]. In [4], the authors investigate in details the relationship between such rings and some homological conjectures, notably the Auslander-Reiten Conjecture and the Gorenstein Symmetry Conjecture etc..

However, it seems there are no enough arguments about the number b_M (called the left Auslander bound of M, see Definition 2.1) for a fixed M in the existing references. The study in this note is a contribution in this direction. We investigate the basic properties of Auslander bounds and give some methods to calculate them (Proposition 2.3 and 2.7, Theorems 2.10 and 2.12). As we show in the paper, the results obtained in this way are also useful to give a new sight of some known theorems and to obtain new results (Corollary 2.11, Propositions 3.7, 3.9 and 3.11).

Since the Auslander Conjecture fails in general even for artin algebras, we naturally consider the (left version of) finitistic Auslander Conjecture as follows.

(fAC) *For every artin algebra* R*, there is an integer* n *such that the left Auslander bound of every finitely generated* R*-module with finite left Auslander bound is not more than* n*.*

We show that the finitistic Auslander bound conjecture for all artin algebras implies the little finitistic dimension conjecture for all artin algebras (Proposition 3.4) and give some cases in which the finitistic Auslander bound conjecture holds (Propositions 3.5).

We now introduce the notions.

Let R be a ring. Given an R-module M and an integer $t \geq 0$, we denote by $M^{>t}$ (${}^{>t}M$, resp.) the subcategory of all modules N such that $\mathrm{Ext}^i_R(M, N) = 0$ ($\mathrm{Ext}^i_R(N, M) = 0$, resp.) for all $i > t$. The notion $M^{\gg 0}$ (${}^{\gg 0}M$, resp.) denotes the subcategory of all modules N such that $N \in M^{>t}$ ($N \in {}^{>t}M$, resp.) for some $t > 0$.

For every R-module M, we denote by $\mathrm{Add}M$ ($\mathrm{add}M$, resp.) the class of all modules isomorphic to direct summands of direct sums (finite direct sums, resp.) of copies of T. We use $\mathrm{pd}M$ ($\mathrm{id}M$, $\mathrm{fd}M$, resp.) to denote the projective (injective, flat, resp.) dimension of M.

In case R is an artin algebra, we denote by $\mathbf{D}$ the usual dual functor between mod$R$ and modR^o. Thus, $(\mathbf{D}(M))_R$ is just the dual module of $M \in \mathrm{mod}R$.

2 Auslander bounds

We prefer to introduce the following notion for the number b_M in the Auslander conjecture.

Definition 2.1 *Let R be a ring and $M \in \mathrm{Mod}R$. The left (right, resp.) Auslander bound of M is defined to be the minimal non-negative integer m such that $M^{\gg 0} = M^{>m}$ (${}^{\gg 0}M = {}^{>m}M$, resp.) , denoted by $\mathrm{lAb}M = m$ ($\mathrm{rAb}M = m$, resp.). If there is no such an integer, we denote that $\mathrm{lAb}M = \infty$ ($\mathrm{rAb}M = \infty$, resp.).*

Let R be a ring and $\mathcal{C}$ be a class of R-modules. We denote by $\mathrm{lAb}|_{\mathcal{C}}M$ the minimal non-negative integer t such that $M^{\gg 0} \bigcap \mathcal{C} = M^{>t} \bigcap \mathcal{C}$. The notion $\mathrm{rAb}|_{\mathcal{C}}M$ is defined similarly. We define $\mathrm{glAb}\mathcal{C} := sup\{\mathrm{lAb}M \mid M \in \mathcal{C}\}$ and $\mathrm{glAb}_{\mathcal{C}}\mathcal{C} := sup\{\mathrm{lAb}_{\mathcal{C}}M \mid M \in \mathcal{C}\}$. The notions $\mathrm{grAb}\mathcal{C}$ and $\mathrm{grAb}_{\mathcal{C}}\mathcal{C}$ are defined similarly.

Remark 2.2 (1) *$\mathrm{lAb}M$ is just the minimal bound on the vanishing of $\mathrm{Ext}_R(M,-)$ in [6]. In [9, 16] etc., $\mathrm{lAb}|_{\mathrm{mod}R}M$ is also denoted by $e_R(M,-)$ in case $M \in \mathrm{mod}R$.*

(2) *One can also define the Tor Auslander bound of M to be the minimal non-negative integer m such that $\mathrm{KerTor}^R_{\gg 0}(M,-) = \mathrm{KerTor}^R_{>m}(M,-)$ (or ∞ if there is no such an integer) and denote by $\mathrm{tAb}M = m$ (or ∞). Then $\mathrm{tAb}M$ is just the minimal bound on the vanishing of $\mathrm{Tor}^R(M,-)$ in [13].*

(3) *Obviously, $\mathrm{lAb}M \leq \mathrm{pd}M$ and $\mathrm{rAb}M \leq \mathrm{id}M$ with the equation holds if the latter are finite. If $M \in \mathrm{mod}R$, then $\mathrm{lAb}|_{\mathrm{mod}R}M = \mathrm{pd}M$ if $\mathrm{pd}M < \infty$.*

Proposition 2.3 $\mathrm{glAb}_{\mathcal{C}}\mathcal{C} = \mathrm{grAb}_{\mathcal{C}}\mathcal{C}$.

Proof. Assume that $\mathrm{grAb}_{\mathcal{C}}\mathcal{C} = t < \infty$. Then for any $M \in \mathcal{C}$ and $N \in M^{\gg 0} \bigcap \mathcal{C}$, we have that $M \in {}^{\gg 0}N \bigcap \mathcal{C} = {}^{>t}N \bigcap \mathcal{C}$ by the definition of $\mathrm{grAb}_{\mathcal{C}}\mathcal{C}$. Hence $N \in M^{>t} \bigcap \mathcal{C}$. It follows that $M^{\gg 0} \bigcap \mathcal{C} = M^{>t} \bigcap \mathcal{C}$. Therefore, $\mathrm{lAb}|_{\mathcal{C}}M \leq t$ and consequently, $\mathrm{glAb}|_{\mathcal{C}}\mathcal{C} \leq t$. Similarly, we have also $\mathrm{grAb}|_{\mathcal{C}}\mathcal{C} \leq \mathrm{glAb}|_{\mathcal{C}}\mathcal{C}$.

#

The above result suggests that we can say the *global Auslander bound* of a class $\mathcal{C}$ of R-modules which is either $\mathrm{grAb}|_{\mathcal{C}}\mathcal{C}$ or $\mathrm{glAb}|_{\mathcal{C}}\mathcal{C}$. This invariance is then denoted by $\mathrm{gAb}\mathcal{C}$.

We denote $\mathrm{GAb}R := \mathrm{gAb}(\mathrm{Mod}R)$ and $\mathrm{gAb}R := \mathrm{gAb}(\mathrm{mod}R)$. Note that $\mathrm{gAb}R$ is just the Ext-index of R in [9, 16] etc.

Proposition 2.4 *Let R be a ring and $\mathcal{C} \subseteq \mathcal{D} \subseteq \mathrm{Mod}R$. Then*

(1) $\mathrm{lAb}|_{\mathcal{C}}M \leq \mathrm{lAb}|_{\mathcal{D}}M$ *and* $\mathrm{rAb}|_{\mathcal{C}}M \leq \mathrm{rAb}|_{\mathcal{D}}M$ *for each* $M \in \mathrm{Mod}R$.

(2) $\mathrm{gAb}\mathcal{C} \leq \mathrm{gAb}\mathcal{D}$.

(3) $\mathrm{glAb}\mathcal{C} \leq \mathrm{glAb}\mathcal{D}$ *and* $\mathrm{grAb}\mathcal{C} \leq \mathrm{grAb}\mathcal{D}$.

(4) $\mathrm{gAb}\mathcal{C} \leq min\{\mathrm{grAb}\mathcal{C}, \mathrm{glAb}\mathcal{C}\}$.

Proof. (1) Assume that $\mathrm{lAb}|_{\mathcal{D}}M = t < \infty$, then $M^{\gg 0} \bigcap \mathcal{D} = M^{>t} \bigcap \mathcal{D}$. Since $\mathcal{C} \subseteq \mathcal{D}$, we have that $M^{\gg 0} \bigcap \mathcal{C} = M^{\gg 0} \bigcap \mathcal{D} \bigcap \mathcal{C} = M^{>t} \bigcap \mathcal{D} \bigcap \mathcal{C} = M^{>t} \bigcap \mathcal{C}$. Hence, $\mathrm{lAb}|_{\mathcal{C}}M \leq t$. The remained part is proved similarly.

(2) Assume that $\mathrm{gAb}\mathcal{D} = t < \infty$, then for any $M \in \mathcal{D}$, it holds $\mathrm{lAb}|_{\mathcal{D}}M \leq t$. As $\mathcal{C} \subseteq \mathcal{D}$, we see that, for any $N \in \mathcal{C}$, $\mathrm{lAb}|_{\mathcal{C}}N \leq \mathrm{lAb}|_{\mathcal{D}}N \leq t$ by (1). It follows that $\mathrm{gAb}\mathcal{C} \leq \mathrm{gAb}\mathcal{D}$.

(3) Assume that $\mathrm{glAb}\mathcal{D} = t < \infty$, then for any $M \in \mathcal{C} \subseteq \mathcal{D}$, it holds $\mathrm{lAb}M \leq t$. therefore, $\mathrm{glAb}\mathcal{C} \leq \mathrm{glAb}\mathcal{D}$. Similarly, $\mathrm{grAb}\mathcal{C} \leq \mathrm{grAb}\mathcal{D}$.

(4) Since $\mathcal{C} \subseteq \mathrm{Mod}R$, $\mathrm{gAb}\mathcal{C} = \mathrm{glAb}|_{\mathcal{C}}\mathcal{C} \leq \mathrm{glAb}\mathcal{C}$ by (1). Similarly, $\mathrm{gAb}\mathcal{C} \leq \mathrm{grAb}\mathcal{C}$, too.

#

Let R be a ring and $\mathcal{C} \subseteq \mathrm{Mod}R$. We say that $\mathcal{C}$ has the two-out-of-three property provided any two terms in a short exact sequence are in $\mathcal{C}$ implies the third term is also in $\mathcal{C}$.

Lemma 2.5 *Let R be a ring and $M \in \mathrm{Mod}R$.*

(1) *All subcategories* $\mathrm{mod}R$, $M^{\gg 0}$, *and* ${}^{\gg 0}M$ *have the two-out-of-three property.*

(2) *All subcategories* $\mathrm{mod}R$, $M^{\gg 0}$,*and* ${}^{\gg 0}M$ *are closed under direct summands and finite direct sums. Moreover,* $M^{\gg 0}$ $({}^{\gg 0}M, resp.)$ *is closed under arbitrary direct sum* $(direct\ products, resp.)$.

(3) $M^{\gg 0} = (\Omega^i M)^{\gg 0}$, *where $\Omega^i M$ denotes an i-th syzygy of M.*

(4) ${}^{\gg 0}M = {}^{\gg 0}(\Omega^{-i}M)$, *where $\Omega^{-i}M$ denotes an i-th cosyzygy of M.*

(5) *If R is an artin algebra and $M \in \mathrm{mod}R$, then* $\mathbf{D}(M^{\gg 0}) = {}^{\gg 0}(\mathbf{D}(M))$.

Proof. Easily.

#

Lemma 2.6 *Let R be a ring and $\mathcal{C} \subseteq \mathrm{Mod}R$. Let $M, N \in \mathrm{Mod}R$.*

(1) $\mathrm{lAb}|_{\mathcal{C}}(M \oplus N) \leq max\{\mathrm{lAb}|_{\mathcal{C}}M, \mathrm{lAb}|_{\mathcal{C}}N\}$.

(2) $\mathrm{rAb}|_{\mathcal{C}}(M \oplus N) \leq max\{\mathrm{rAb}|_{\mathcal{C}}M, \mathrm{rAb}|_{\mathcal{C}}N\}$.

(3) *If R is an artin algebra and $M \in \mathrm{mod}R$, then* $\mathrm{lAb}|_{\mathrm{mod}R}M = \mathrm{rAb}|_{\mathrm{mod}R}(\mathbf{D}(M))_R$.

Proof. (1) Clearly we can assume that $k = max\{\mathrm{lAb}|_{\mathcal{C}}M, \mathrm{lAb}|_{\mathcal{C}}N\} < \infty$. Now note that $(M \oplus N)^{\gg 0} \bigcap \mathcal{C} = M^{\gg 0} \bigcap N^{\gg 0} \bigcap \mathcal{C} = M^{>k} \bigcap N^{>k} \bigcap \mathcal{C} = (M \oplus N)^{>k} \bigcap \mathcal{C}$, so we have that $\mathrm{lAb}|_{\mathcal{C}}(M \oplus N) \leq k$.

(2) Dually to (1).

(3) By Lemma 2.5 and the definitons.

#

Proposition 2.7 *Let R be a ring and $\mathcal{C} \subseteq \mathrm{Mod}R$. Let $M, N \in \mathrm{Mod}R$.*

(1) $\mathrm{lAb}|_{\mathcal{C}}M \leq m$ *if and only if* $\mathrm{lAb}|_{\mathcal{C}}(\Omega^m M) = 0$, *where $\Omega^m M$ denotes an m-th syzygy of M.*

(2) $\mathrm{rAb}|_{\mathcal{C}}M \leq m$ *if and only if* $\mathrm{rAb}|_{\mathcal{C}}(\Omega^{-m} M) = 0$, *where $\Omega^{-m} M$ denotes an m-th cosyzygy of M.*

Proof. We still prove (1) and left (2) to the readers.

Since $M^{\gg 0} = (\Omega^m M)^{\gg 0}$ by Lemma 2.5 and $M^{>m} = (\Omega^m M)^{\geq 1}$ by the dimension shifting, we see that $\mathrm{lAb}|_{\mathcal{C}}M \leq m \iff M^{\gg 0} \bigcap \mathcal{C} \subseteq M^{>m} \bigcap \mathcal{C} \iff (\Omega^m M)^{\gg 0} \bigcap \mathcal{C} \subseteq (\Omega^m M)^{\geq 1} \bigcap \mathcal{C} \iff \mathrm{lAb}|_{\mathcal{C}}(\Omega^m M) = 0$.

#

The following result is now obvious.

Proposition 2.8 *The following are equivalent for the ring R and $\mathcal{C} \subseteq \mathrm{Mod}R$.*

(1) $\mathrm{gAb}\mathcal{C} \leq n$

(2) $\mathrm{Ext}_R^{i>n}(M, N) = 0$ *for all $M, N \in \mathcal{C}$ such that* $\mathrm{Ext}_R^i(M, N) = 0$ *for all sufficiently large i.*

(3) $\mathrm{glAb}|_{\mathcal{C}}(\Omega^n \mathcal{C}) = 0$, *where $\Omega^n \mathcal{C}$ denotes the class of all n-th syzygies of R-modules in $\mathcal{C}$.*

(4) $\mathrm{grAb}|_{\mathcal{C}}(\Omega^{-n} \mathcal{C}) = 0$, *where $\Omega^{-n} \mathcal{C}$ denotes the class of all n-th cosyzygies of R-modules in $\mathcal{C}$.*

Proposition 2.9 *Let R be an artin algebra. Then* $\mathrm{gAb}R = \mathrm{gAb}R^o$.

Proof. Since R is an artin algebra, there is a duality $\mathbf{D}$ between $\mathrm{mod}R$ and $\mathrm{mod}R^o$. Thus, we obtain that $\mathrm{gAb}R \leq n$ if and only if $\mathrm{Ext}_R^{i>n}(M, N) = 0$ for all $M, N \in \mathrm{mod}R$ such that $\mathrm{Ext}_R^i(M, N) = 0$ for all sufficiently large i, if and only if $\mathrm{Ext}_R^{i>n}(\mathbf{D}M, \mathbf{D}N) = 0$ if and only if $\mathrm{gAb}R^o \leq n$.

#

Theorem 2.10 *Let R be a ring and $M \in \mathrm{Mod}R$.*

(1) *If* $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty$ *and* $R \in M^{\gg 0}$, *then* $\mathrm{lAb}|_{\mathrm{mod}R}M = min\{t | R \in M^{>t}\}$.

(2) *If* $\mathrm{lAb}M < \infty$ *and* $R^{(\kappa)} \in M^{\gg 0}$ *for all cardinals κ, then* $\mathrm{lAb}M = min\{t | R^{(\kappa)} \in M^{>t}$ *for all cardinals* $\kappa\}$.

(3) *If* $\mathrm{rAb}M < \infty$ *and* $I \in {}^{\gg 0}M$ *for all injective* R*-module* I*, then* $\mathrm{rAb}M = min\{t | I \in {}^{>t}M$ *for all injective* R*-module* $I\}$.

Proof. (1) Assume that $\mathrm{lAb}|_{\mathrm{mod}R}M = m < \infty$ and $t = min\{t | R \in M^{>t}\} < \infty$.

On the one hand, since $R \in M^{\gg 0} \bigcap \mathrm{mod}R$, $R \in M^{>m}$ by the definition of $\mathrm{lAb}|_{\mathrm{mod}R}M$. It follows that $t \leq m$.

On the other hand, by the definition of t, we see that $R \in M^{>t} \bigcap \mathrm{mod}R \subseteq M^{\gg 0} \bigcap \mathrm{mod}R$. Now take any $N \in M^{\gg 0} \bigcap \mathrm{mod}R$ and any projective resolution of N: $\cdots \to P_1 \to P_0 \to N \to 0$ with each $P_i \in \mathrm{mod}R$ finitely generated projective. Then all $P_i \in M^{>t} \bigcap \mathrm{mod}R$, and also all $\Omega^i N \in M^{\gg 0} \bigcap \mathrm{mod}R$, by Lemma 2.5. Therefore, for all $i > t$, we obtain that $\mathrm{Ext}^i_R(M, N) \simeq \mathrm{Ext}^{i+1}_R(M, \Omega N) \simeq \cdots \simeq \mathrm{Ext}^{i+m}_R(M, \Omega^m N) = 0$, by the dimension shifting and the definition of $\mathrm{lAb}|_{\mathrm{mod}R}M$. It follows that $N \in M^{>t} \bigcap \mathrm{mod}R$. Consequently, $M^{>t} \bigcap \mathrm{mod}R = M^{\gg 0} \bigcap \mathrm{mod}R$, that is, $\mathrm{lAb}|_{\mathrm{mod}R}M \leq t$.

The proof of (2) is similar as (1) and the proof of (3) is dual to (2).

#

The following result is now a direct corollary of the above theorem (see for instance [16, Corollary 3.3]).

Corollary 2.11 *Let* R *be a ring with* $\mathrm{id}R < \infty$*.*

(1) *If* $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty$ *for every* $M \in \mathrm{mod}R$*, then* $\mathrm{gAb}R = \mathrm{id}R$*.*

(2) *If* R *is left noetherian and* $\mathrm{lAb}M < \infty$ *for every* $M \in \mathrm{Mod}R$*, then* $\mathrm{GAb}R = \mathrm{id}R$*.*

We note that assumptions in Theorem 2.10 can not be removed. As to the first example, let R be an artin algebra of finite representation type with $\mathrm{id}R = \infty$. Then it is easy to see that $\mathrm{gAb}R < \infty$. However, it is clear that there are modules $M \in \mathrm{mod}R$ such that $min\{t | R \in M^{>t}\} = \infty$. To the second example, let R be a Gorenstein ring with $\mathrm{gAb}R = \infty$ (such rings exist by [11]). Then there are modules $M \in \mathrm{mod}R$ such that $\mathrm{lAb}|_{\mathrm{mod}R}M = \infty$ by Corollary 2.11. However, it is easy to see that $min\{t | R \in M^{>t}\} \leq \mathrm{id}R < \infty$ for any $M \in \mathrm{mod}R$.

It is a question whether $\mathrm{GAb}R = \mathrm{gAb}R$ if R is left noetherian.

Theorem 2.12 *Let* R *be a ring and* $M \in \mathrm{Mod}R$*.*

(1) *Assume that* $M \oplus R \in M^{\gg 0} \bigcap \mathrm{mod}R$*, then* $\mathrm{lAb}|_{\mathrm{mod}R}M = \mathrm{pd}M$*.*

(2) *Assume that* $M \oplus R^{(\kappa)} \in M^{\gg 0}$ *for any cardinal* κ*, then* $\mathrm{lAb}M = \mathrm{pd}M$*.*

(3) *Assume that* $M, I \in {}^{\gg 0}M$ *for any injective* R*-module* I*, then* $\mathrm{rAb}M = \mathrm{id}M$*.*

Proof. (1) Clearly we need only prove that $\mathrm{pd}M \leq \mathrm{lAb}_{\mathrm{mod}R}M$.

We can assume that $\mathrm{lAb}_{\mathrm{mod}R}M = m < \infty$. Since $M \in \mathrm{mod}R$, we can take a projective resolution of M: $\cdots \to P_1 \to P_0 \to M \to 0$ with each P_i finitely generated projective. Then each $\Omega^i M \in M^{\gg 0} \bigcap \mathrm{mod}R$ by Lemma 2.5, as $M \oplus R \in M^{\gg 0}$. It follows that $\Omega^i M \in M^{>m}$ for each i, by the definition of $\mathrm{lAb}_{\mathrm{mod}R}M$. Now by applying the functor $\mathrm{Hom}_R(-, \Omega^{m+1}M)$ to the exact sequence $0 \to \Omega^m M \to P_{m-1} \to \cdots \to P_0 \to M \to 0$, we obtain that $\mathrm{Ext}^1_R(\Omega^m M, \Omega^{m+1}M) \simeq \mathrm{Ext}^2_R(\Omega^{m-1}M, \Omega^{m+1}M) \simeq \cdots \simeq \mathrm{Ext}^{m+1}_R(M, \Omega^{m+1}M) = 0$ by the dimension shifting. It follows that the exact sequence $0 \to \Omega^{m+1}M \to P_m \to \Omega^m M \to 0$ splits, and consequently, $\mathrm{pd}M \leq m$.

(2) Similarly as (1).

(3) Dually to (2).

#

3 Homological conjectures

Let R be a ring. Denote $\mathrm{fPD}(R) := sup\{\mathrm{pd}M \mid \mathrm{pd}M < \infty \text{ and } M \in \mathrm{mod}R\}$ and $\mathrm{FPD}(R) := sup\{\mathrm{pd}M \mid \mathrm{pd}M < \infty\}$. Similarly, we denote $\mathrm{fID}(R) := sup\{\mathrm{id}M \mid \mathrm{id}M < \infty \text{ and } M \in \mathrm{mod}R\}$ and $\mathrm{FID}(R) := sup\{\mathrm{id}M \mid \mathrm{id}M < \infty\}$.

Recall first the following open conjecture.

Finitistic Dimension Conjecture (fDC):

(fPDC) $\mathrm{fPD}(R) < \infty$ *for every artin algebra* R*.* Or dually

(fIDC) $\mathrm{fID}(R) < \infty$ *for every artin algebra* R*.*

It is clear that (fPDC) holds for an artin algebra R if and only if (fIDC) holds for R^o.

The following result gives a relationship between the finitistic dimension and the Auslander bound.

Proposition 3.1 *Let* R *be a ring.*

(1) $\mathrm{fPD}(R) \leq \mathrm{rAb}|_{\mathrm{mod}R}R$.

(2) *If* R *is an artin algebra, then* $\mathrm{fID}(R) \leq \mathrm{lAb}|_{\mathrm{mod}R}\mathbf{D}(R_R)$.

Proof. (1) Let $M \in \mathrm{mod}R$. Since $\mathrm{pd}M \leq n \Longleftrightarrow \mathrm{Ext}^{i>n}_R(M, R) = 0$ and $\mathrm{pd}M < \infty$ (see for instance [7, Theorem 3], the conclusion follows from the definition of $\mathrm{rAb}|_{\mathrm{mod}R}R$.

(2) By (1) and the duality in the artin algebras.

#

The following result can also be seen from Proposition 3.1. We give here another proof.

Proposition 3.2 *Let* R *be an artin algebra. If* $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty$ ($\mathrm{rAb}|_{\mathrm{mod}R}M < \infty$*, resp.*)*, for every* $M \in \mathrm{mod}R$*, then* (fIDC)((fPDC)*, resp.*) *holds for* R*.*

Proof. Since R is an artin algebra, every module has finite length. Let $\mathcal{S} = \{S_1, \cdots, S_n\}$ be the set of all simple modules. Assume that $\mathrm{id}M < \infty$. Then $M \in (\oplus S_i)^{\gg 0} = (\oplus S_i)^{> \mathrm{lAb}(\oplus S_i)}$. It follows that $\mathrm{id}M \leq \mathrm{lAb}(\oplus S_i) < \infty$ by assumptions. The remained part can be proved similarly.

#

Let R be a finite dimensional algebra over a field and let R^e denote the enveloping algebra of R. There is a relation between the finitistic projective dimension of R and the left Auslander bound of R as an R^e-module.

Proposition 3.3 $\mathrm{fPD}(R) \leq \mathrm{lAb}({}_{R^e}R)$.

Proof. See for instance [8, 1 Proposition].

#

As pointed in [11, 18], the Auslander conjecture (stated in the introduction) fails for artin algebras in general. However, we can consider a finitistic version of the Auslander conjecture. Let R be a ring. We denote $\mathrm{FLAb}(R) := sup\{\mathrm{lAb}M \mid \mathrm{lAb}M < \infty\}$ and $\mathrm{fLAb}(R) := sup\{\mathrm{lAb}|_{\mathrm{mod}R}M \mid M \in \mathrm{mod}R$. Similarly, $\mathrm{FRAb}(R) = sup\{\mathrm{rAb}M \mid \mathrm{rAb}M < \infty\}$ and $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty\}$ and $\mathrm{fRAb}(R) = sup\{\mathrm{rAb}|_{\mathrm{mod}R}M \mid M \in \mathrm{mod}R$ and $\mathrm{rAb}|_{\mathrm{mod}R}M < \infty\}$.

Finitistic Auslander Conjecture (FAC):

(lFAC) $\mathrm{fLAb}(R) < \infty$ *for every artin algebra* R. Or dually

(rFAC) $\mathrm{fRAb}(R) < \infty$ *for every artin algebra* R.

It is clear that $\mathrm{fLAb}R = \mathrm{fRAb}R < \infty$ if $\mathrm{gAb}R < \infty$ and that $\mathrm{FLAb}R = \mathrm{FRAb}R < \infty$ if $\mathrm{GAb}R < \infty$ by Proposition 2.3. Note also that $\mathrm{fLAb}(R) = \mathrm{fRAb}(R^o)$ in case that R is an artin algebra.

It was noticed that every group algebra kG with k a field and G finite has the property $\mathrm{GAb}(kG) < \infty$, see [3, Theorem 2.4] and [4, Appendix A].

It is also interesting to know whether $\mathrm{fLAb}(R) = \mathrm{FLAb}(R)$ for every artin algebra R.

Proposition 3.4 *Let* R *be a ring.*

(1) $\mathrm{fLAb}(R) < \infty$ *implies* $\mathrm{fPD}(R) < \infty$.

(1′) $\mathrm{FLAb}(R) < \infty$ *implies* $\mathrm{FPD}(R) < \infty$.

(2) $\mathrm{fRAb}(R) < \infty$ *implies* $\mathrm{fID}(R) < \infty$.

(2′) $\mathrm{FRAb}(R) < \infty$ *implies* $\mathrm{FID}(R) < \infty$.

Proof. Directly by Remark 2.2 (3).

#

As (fPDC) fails for commutative noetherian rings in general, (FAC) fails too in the case. Moreover, it is pointed that there is a commutative notherian ring R with infinite Krull dimension such that $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty$ for every $M \in \mathrm{mod}R$ but $\mathrm{gAb}R = \infty$ (c.f. [4, Remark 3.3]). However, it is interesting to check (FAC) in case of artin algebras. In particular, it is not known whether $\mathrm{gAb}R < \infty$ if R is an artin algebra such that $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty$ for every $M \in \mathrm{mod}R$.

The following result gives partial answer to (FAC).

Proposition 3.5 *Let R be a ring.*

(1) *If* $\mathrm{id}R < \infty$*, then* $\mathrm{lFAb}(R) < \infty$*.*

(2) *If R is left noetherian and* $\mathrm{id}R < \infty$*, then* $\mathrm{LFAb}(R) < \infty$*.*

Proof. (1) Indeed, we have that $\mathrm{lAb}|_{\mathrm{mod}R}M = min\{t|R \in M^{>t}\} \leq \mathrm{id}R$ provided $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty$ and $M \in \mathrm{mod}R$, by Theorem 2.10.

(2) If R is left noetherian and $\mathrm{id}R < \infty$, then $\mathrm{id}R^{(\kappa)} < \infty$ for any cardinal κ. Now the conclusion follows from Theorem 2.10 again.

#

Let R be a ring and $T \in \mathrm{Mod}R$ with $S = \mathrm{End}_R T$. Recall from [19] that T is Wakamatsu-tilting if it satisfies (1) $T \in \mathrm{mod}R$ and $T_S \in \mathrm{mod}S^o$, (2) $R \simeq \mathrm{End}(T_S)$, and (3) $\mathrm{Ext}^i_R(T,T) = 0 = \mathrm{Ext}^i_{S^o}(T,T)$ for all $i > 0$. Equivalently, T is Wakamatsu-tilting if (W1) $T \in \mathrm{mod}R$, (W2) $\mathrm{Ext}^i_R(T,T) = 0$ for all $i > 0$, and (W3) there is an exact sequence $0 \to R \to^{f_0} T_0 \to^{f_1} T_1 \to^{f_2} \cdots$ with each $T_i \in \mathrm{add}T$ and each $\mathrm{Im}f_i \in {}^{>0}T$. It is clear that T is Wakamatsu-tilting if and only if T_S is Wakamatsu-tilting. Recall also that T is tilting if it satisfies (T1) $T \in \mathrm{mod}R$ and $\mathrm{pd}T < \infty$, (T2) $\mathrm{Ext}^i_R(T,T) = 0$ for all $i > 0$, and (T3) there is an exact sequence $0 \to R \to T_0 \to \cdots \to T_n \to 0$ with each $T_i \in \mathrm{add}T$, for some integer n. We note that T is tilting if and only if T_S is tilting, where $S = \mathrm{End}_R T$ [15].

Wakamatsu Tilting Conjecture: (WTC)

(PWTC) *Every Wakamatsu-tilting module of finite projective dimension is tilting.*

It is proved in [14] that, if $\mathrm{fPD}(R) < \infty$ for a ring R, then (PWTC) holds for R.

Proposition 3.6 (PWTC) *holds for all rings if and only if the following conjecture holds for all rings.*

(EWTC) *An R-module $T \in \mathrm{mod}R$ is tilting if it satisfies the conditions* (T2) *and* (T3) *in the definition of tilting modules.*

Proof. Let R be a ring and $T \in \mathrm{mod}R$ with $S = \mathrm{End}_R T$.

(EWTC) ⇒ (PWTC): Assume that T is Wakamatsu-tilting with $\mathrm{pd}T < \infty$. Then we have an exact sequence $0 \to P_n \to \cdots \to P_0 \to T \to 0$ for some n. Applying the functor $\mathrm{Hom}_R(-,T)$, we obtain an induced exact sequence $0 \to S \to T_0 \to \cdots \to T_n \to 0$ since $\mathrm{Ext}^i_R(T,T) = 0$ for all $i > 0$. Note that $\mathrm{Ext}^i_{S^o}(T,T) = 0$ for all $i > 0$ since T is Wakamatsu-tilting, we get that T is tilting if (EWTC) holds for S^o.

(PWTC) $\Rightarrow$ (EWTC): If (T2) and (T3) in the definition of tilting modules holds for $T \in \mathrm{mod}R$, then T is Wakamatsu-tilting. Applying the functor $\mathrm{Hom}_R(-,T)$ to the exact sequence in (T3), we easily see that T_S is Wakamatsu-tilting with finite projective dimension. It follows that T_S is tilting if (PWTC) holds for S^o. Now by the left-right symmetry, we get that T is tilting.

#

If we specify $n = 0$ in the condition (T3), then $R \in \mathrm{add}T$. In this case, the conjecture (EWTC) is just the Auslander-Reiten conjecture.

Auslander-Reiten Conjecture: (ARC) *Every selforthogonal generator possessing finitely generated projective resolutions is projective.*

The following result gives a partial answer to (EWTC), which extends [4, Theorem A].

Proposition 3.7 *Let R be a ring. If $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty$ for every $M \in \mathrm{mod}R$, then the conjecture* (EWTC) *holds for R. In particular,* (ARC) *holds for R.*

Proof. Assume that $T \in \mathrm{mod}R$ satisfies the conditions (T2) and (T3). Then we easily obtain that $T \oplus R \in T^{\gg 0}$. Now by the assumption and Proposition 2.12, we get that $\mathrm{pd}T = \mathrm{lAb}|_{\mathrm{mod}R}T < \infty$. It follows that T is tilting by the definition.

#

The conjecture (EWTC) has an infinitely generated version. Let R be a ring and $T \in \mathrm{Mod}R$. Recall from [1] that T is infinitely generated tilting if it satisfies (T1') $\mathrm{pd}T < \infty$, (T2') $\mathrm{Ext}^i_R(T, T^{(\kappa)}) = 0$ for all cardinals κ and all $i > 0$, and (T3') there is an exact sequence $0 \to R \to T_0 \to \cdots \to T_n \to 0$ with each $T_i \in \mathrm{Add}T$. Similarly as the above result, we also have the following.

Proposition 3.8 *Let R be a ring and $T \in \mathrm{Mod}R$. If $\mathrm{lAb}M < \infty$ for every $M \in \mathrm{Mod}R$, then T is infinitely generated tilting if and only if it satisfies* (T2$'$) *and* (T3$'$) *in the definiton. In particular, every selforthogonal generator is projective.*

Proof. Similarly as Proposition 3.7.

#

Another special case of (WTC) is as follows (indeed, restricting to the trivial Wakamatsu-tilting module $\mathbf{D}R$).

Gorenstein Symmetry Conjecture (GSC): *Let R be an artin algebra. Then* $\mathrm{id}R < \infty$ *if and only if* $\mathrm{id}(R_R) < \infty$.

(GSC) clearly makes sense for any ring. It was proves in [4] that if R is a two-sided noetherian ring such that $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty$ for every $M \in \mathrm{mod}R$ and $\mathrm{lAb}|_{\mathrm{mod}R^o}M < \infty$ for every $M \in \mathrm{mod}R^o$, and (1) R is an artin algebra, or (2) R has a dualizing complex, then $\mathrm{id}R < \infty$ if and only if $\mathrm{id}_{R^o}R < \infty$ (whence, $\mathrm{id}R = \mathrm{id}_{R^o}R$ by [10]). The following result gives another answer. Note that we do not know whether $\mathrm{lAb}M < \infty$ for every $M \in \mathrm{Mod}R$ if $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty$ for every $M \in \mathrm{mod}R$, even when R is an artin algebra. Yet we do not know whether $\mathrm{lAb}M < \infty$ for every $M \in \mathrm{Mod}R$ implies that $\mathrm{rAb}M < \infty$ for every $M \in \mathrm{Mod}R$.

Proposition 3.9 *Let R be a two-sided noetherian ring. Assume that*

(1) $\mathrm{lAb}M < \infty$ *for every $M \in \mathrm{Mod}R$ and every $M \in \mathrm{Mod}R^o$, or*

(2) *R has a dualizing complex and* $\mathrm{rAb}M < \infty$ *for every $M \in \mathrm{mod}R$ and every $M \in \mathrm{mod}R^o$.*

Then $\mathrm{id}R < \infty$ *if and only if* $\mathrm{id}R^o < \infty$.

Proof. Assume (1) holds. If $\mathrm{id}R < \infty$, then $\mathrm{GAb}R < \infty$ by assumptions and Corollary 2.11. It follows from Proposition 3.4 that $\mathrm{FID}(R) < \infty$. Hence, $\mathrm{id}R^o < \infty$ by [7, Proposition 7]. Another part is symmetry.

Assume now (2) holds. If $\mathrm{id}R < \infty$, then for any $M \in \mathrm{mod}R$ and any injective R^o-module N, it holds that $\mathrm{Tor}^R_{(\mathrm{id}R)+1}(N, M) \simeq \mathrm{Hom}_{R^o}(\mathrm{Ext}_R^{(\mathrm{id}R)+1}(M, R), N) = 0$. Hence $\mathrm{fd}_{R^o}N < \infty$. Since R^o has a dualizing complex, all R^o-modules of finite flat dimension have finite projective dimension, by [12]. It follows that $N \in {}^{\gg 0}_{R^o}R$ for any injective R^o-module N. Now applying Theorem 2.12 to the R^o-module R, we obtain that $\mathrm{id}_{R^o}R = \mathrm{rAb}_{R^o}R < \infty$ by assumptions. The other part is symmetry.

#

Let R be a ring. Recall that an R-module M has a projective resolution with a strongly redundant image from m if the m-th syzygy $\Omega^m M \simeq \oplus M_i$ with each $M_i \in \mathrm{add}(\Omega^{m_i}M)$ for some $m_i > m$. The dual concept pertains to injective resolutions. According to [7, Lemma 4] and its dual, for each $i \geq m$, every direct summand of $\Omega^i M$ ($\Omega^{-i}M$, resp.) appears as a direct summand of some syzygies (cosyzygies, resp.) latter than i in case M has a projective (injective, resp.) resolution with a strongly redundant image from m.

Lemma 3.10 *Let R be a ring. If $M \in \mathrm{mod}R$ has a projective resolution with a strongly redundant image from some integer m, then* $\mathrm{lAb}|_{\mathrm{mod}R}M = m < \infty$. *Dually, if $M \in \mathrm{mod}R$ has a injective resolution with a strongly redundant image from some integer m, then* $\mathrm{rAb}|_{\mathrm{mod}R}M = m < \infty$.

Proof. Since M has a projective resolution with a strongly redundant image from m, we have that, for every $i \geq m$, each direct summand of $\Omega^i M$ appears in the latter syzygies. Now take any $N \in M^{\gg 0} \bigcap \mathrm{mod}R$ and assume that $N \in M^{>t}$. It follows that $\mathrm{Ext}^1_R(\Omega^m M, N) \leq \mathrm{Ext}^1_R(\oplus_{i>t}\Omega^i M, N) \simeq \prod_{i>t} \mathrm{Ext}^{i+1}_R(M, N) = 0$ by the dimension shifting. Note that $M^{\gg 0} = (\Omega^m M)^{\gg 0}$ by Lemma 2.5, so $\mathrm{lAb}(\Omega^m M) = 0$. It follows that $\mathrm{lAb}M \leq m$ by Proposition 2.7.

The remained part is proved dually.

#

Let R be a ring. Recall that an R-module M has a ultimately closed projective resolution if the m-th syzygy $\Omega^m M \simeq \oplus M_i$ with each $M_i \in \mathrm{add}(\Omega^{m_i} M)$ for some $m_i < m$. The dual concept also pertains to injective resolutions. One call the rings such that every finitely generated module has a ultimately closed injective (projective, resp.) resolution to be of injective (projective, resp.) ultimately closed type. Obviously, if R is an artin algebra of injective ultimately closed type then R^o is of projective ultimately closed type. It was shown in [17, Section 8] that if R is semi-primary with $R/\mathrm{Rad}^m(R)$ of finite representation type and $\mathrm{Rad}^{m+1}(R) = 0$ then R is of injective ultimately closed type. In fact, the proof given there even shows there is a common integer k such that every module has a ultimately closed injective resolution at k. In particular, semi-primary rings with $\mathrm{Rad}^2 = 0$ have the property.

It was proved in [2] that the Auslander-Reiten Conjecture holds for artin algebras of projective ultimately closed type.

Proposition 3.11 (1) *If R is an artin algebra of projective* (*injective, resp.*) *ultimately closed type, then* $\mathrm{lAb}|_{\mathrm{mod}R}M < \infty$ ($\mathrm{rAb}|_{\mathrm{mod}R}M < \infty$, *resp.*) *for every* $M \in \mathrm{mod}R$. *In particular,* (EWTC) *holds for all artin algebras of projective ultimately closed type.*

(2) *If R is an artin algebra with $R/\mathrm{Rad}^m(R)$ of finite representation type and* $\mathrm{Rad}^{m+1}(R) = 0$*, then* $\mathrm{gAb}R < \infty$.

Proof. It was remarked in [5] that, over an artin algebra, a ultimately closed injective (projective) resolution of a finitely generated module is strongly redundant. Thus the conclusions follow from Lemma 3.10 and Proposition 3.7.

#

The above result also adds artin algebras of projective ultimately closed type to the class of rings satisfying Auslander conjecture.

Results in this section suggest the following conjectures which generalize the Auslander-Reiten Conjecture.

Generalized Auslander-Reiten Conjecture (GARC): Let R be a ring and $M \in \mathrm{mod}R$. If $M \oplus R \in M^{\gg 0}$, then $\mathrm{pd}M < \infty$.

Conjecture B: Let R be a ring and $M \in \mathrm{mod}R$. If $M \in M^{\gg 0}$, then $\mathrm{lAb}M < \infty$.

By Theorem 2.12, Conjecture B implies (GARC). Thus, by Proposition 3.11, both conjectures hold for artin algebras of projective ultimately closed type.

References

[1] Angeleri Hügel L. and Coelho F.U., Infinitely generated tilting modules of finite projective dimension, Forum Math. 13 (2001), 239-250.

[2] Auslander M. and Reiten I., Applications of contravariantly finite subcategories, Adv. Math. 86 (1) (1991), 111-152.

[3] Benson D. J., Carlson J. F., and Robinson G. R., On the vanishing of group cohomology, J. Algebra 131 (1) (1990), 40C73.

[4] Christensen L. W. and Holm H., Algebras that satisfy auslanders condition on vanishing of cohomology, arXiv: 0801.0401v1.

[5] Colby R.R. and Fuller K.R., A note on the Nakayama conjectures, Tsukuba J. Math. 14(2) (1990), 343-352.

[6] Cowley K. M., One-Sided Bounds and the Vanishing of Ext, J. Algebra 190 (1997), 361-371.

[7] Fuller K.R. and Wang Y., Redundancy in resolutions and finitistic dimensions of noetherian rings, Comm. Algebra 21 (8) (1993), 2983-2994.

[8] Happel D., Homological conjectures in representation theory of finite- dimensional algebras, Sherbrook Lecture Notes Series (1991), available from http://www.math.ntnu.no/oyvinso/Nordfjordeid/Program/references.html.

[9] Huneke C. and Jorgensoen D.A., Symmetry in the vanishing of Ext over Gorenstein rings, Math. Scand 93 (8) (2003), 161-184.

[10] Iwanaga Y., On rings with finite self-injective dimension II, Tsukuba J. Math. 4 (1) (1980), 107-113.

[11] Jorgensen D.A. and Sega L.M., Nonvanishing cohomology and classes of Gorenstein rings, Adv. Math. 188 (2) (2004), 470-490.

[12] Jorgensen P., Finite flat and projective dimensions, Comm. Algebra 33 (2005), 2275-2279.

[13] Kirkman E. and Kuzmanovich J., On the finitistic dimension of fixed subrings, Comm. Algebra 22 (1994), 4621-4635.

[14] Mantese F. and Reiten I., Wakamatsu Tilting modules, J. Algebra 278 (2004), 532-552.

[15] Miyashita Y., Tilting modules of finite projective dimension, Math. Zeit. 193 (1986), 113-146.

[16] Mori I., Symmetry in the vanishing of Ext over stably symmetric algebras, J. Algebra 310 (2) (2007), 708-729.

[17] Tachikawa H., Quasi-Frobenius rings and generalizations. QF-3 and QF-1 rings, Springer-Verlag, Berlin, 1973, Notes by Claus Michael Ringel, Lect. Notes Math., Vol. 351.

[18] Smalo S.O., Local limitations of the Ext functor do not exist, Bull. London Math. Soc. 38 (1) (2006), 97-98.

[19] Wakamatsu T., On modules with trivial self-extensions, J. Algebra 114 (1988) 106-114.

[20] Wakamatsu T., Stable equivalence of self-injective algebras and a generalization of tilting modules, J. Algebra 134 (1990), 298-325.